

New Partition Properties under AD

S. Jackson

Department of Mathematics
University of North Texas
jackson@unt.edu

ESI Workshop
June, 2009, Vienna

We work in the theory $\text{ZF} + \text{DC} + \text{AD}$.

Recall the Erdős-Rado Partition notation.

Definition

$\kappa \rightarrow (\kappa)^\lambda$ if for every partition $\mathcal{P}: (\kappa)^\lambda \rightarrow \{0, 1\}$ of the increasing functions from λ to κ into two pieces, there is a homogeneous $H \subseteq \kappa$ of size κ .

We say κ has the **strong** partition property if $\kappa \rightarrow (\kappa)^\kappa$, and say κ has the **weak** partition property if $\kappa \rightarrow (\kappa)^\lambda$ for all $\lambda < \kappa$.

We usually use a reformulation of the partition property which uses c.u.b. homogeneous sets.

To state this, we need the notion of **type** of a function.

Definition

We say $f: \alpha \rightarrow \text{On}$ has **uniform cofinality** ω if there is an $f': \alpha \times \omega \rightarrow \text{On}$ which is increasing in the second argument and $f(\beta) = \sup_n f'(\beta, n)$ for all $\beta < \alpha$.

Definition

$f: \alpha \rightarrow \text{On}$ is of the **correct type** if f is increasing, everywhere discontinuous, and of uniform cofinality ω .

We can likewise define uniform cofinality ω_1, ω_2 , etc.

General Types

More generally, for any $g: \alpha \rightarrow \text{On}$ we can define f having **uniform cofinality g** : there is an

$$f' : \{(\beta, \gamma) : \beta < \alpha, \gamma < g(\beta)\}$$

with $f(\beta) = \sup_{\gamma} f'(\beta, \gamma)$.

We use frequently the “almost everywhere” versions of these notions with respect to some measure μ on $\text{dom}(f)$.

Partition Relations

Definition

$\kappa \xrightarrow{\text{c.u.b.}} (\kappa)^\lambda$ iff for every partition \mathcal{P} of the function $f: \lambda \rightarrow \kappa$ of the correct type, there is a c.u.b. $C \subseteq \kappa$ which is homogeneous for \mathcal{P} .

The ordinary and c.u.b. version of the partition relation are essentially equivalent:

Fact

$$\kappa \xrightarrow{\text{c.u.b.}} (\kappa)^\lambda \Rightarrow \kappa \rightarrow (\kappa)^\lambda$$

$$\kappa \rightarrow (\kappa)^{\omega \cdot \lambda} \Rightarrow \kappa \xrightarrow{\text{c.u.b.}} (\kappa)^\lambda$$

Definition

$A \subseteq \omega^\omega$ is κ -Suslin if there is a tree $T \subseteq \omega \times \kappa$ such that $A = p[T] = \{x : \exists f \in \kappa^\omega \in (x, f) \in [T]\}$. Let $S(\kappa) =$ the κ -Suslin sets.

κ is a Suslin cardinal if $S(\kappa) - \bigcup_{\lambda < \kappa} S(\lambda) \neq \emptyset$.

Theorem (Steel-Woodin)

The Suslin cardinals are closed below their supremum.

Assuming $V = L(\mathbb{R})$, there is a largest Suslin cardinal δ_1^2 (Martin-Steel), and the Suslin cardinals are c.u.b. in δ_1^2 .

The first ω many Suslin cardinals are:

$$\lambda_1 = \omega, \delta_1^1 = \omega_1, \lambda_3 = \omega_\omega, \delta_3^1 = \omega_{\omega^\omega+1}, \lambda_5 = \omega_{\omega^{\omega^\omega}}, \delta_5^1 = \omega_{\omega^{\omega^\omega+1}}, \\ \dots, \lambda_{2n+1}, \delta_{2n+1}^1, \dots$$

Here $\delta_{2n+1}^1 = (\lambda_{2n+1})^+$, and λ_{2n+1} is a cardinal of cofinality ω (Kechris).

Recall $\delta_{2n+2}^1 = (\delta_{2n+1}^1)^+$, and all the δ_n^1 are regular (measurable).

Definition

Given an increasing, discontinuous sequence of cardinals $\{\kappa_i\}_{i<\theta}$ and a sequence of ordinals $\{\lambda_i\}_{i<\theta}$, we say a **block function** from $\vec{\lambda}$ to $\vec{\kappa}$ is a $\vec{f} = \{f_i\}_{i<\theta}$ where $f_i: \lambda_i \rightarrow \kappa_i - \sup_{j<i} \kappa_j$. A block c.u.b. set is a $\vec{C} = \{C_i\}_{i<\theta}$ where C_i is c.u.b. in $\kappa_i - \sup_{j<i} \kappa_j$.

We next define the **polarized** partition property.

Definition

$\vec{\kappa} \rightarrow (\vec{\kappa})^{\vec{\lambda}}$ if for every partition \mathcal{P} of the block functions $\vec{f}: \vec{\lambda} \rightarrow \vec{\kappa}$ into two pieces, there is a block c.u.b. set \vec{C} homogeneous for \mathcal{P} .

For sequences of length 3 we will write

$$(\kappa_0, \kappa_1, \kappa_2) \rightarrow (\kappa_0, \kappa_1, \kappa_2)^{\lambda_0, \lambda_1, \lambda_2}.$$

Main Results

Theorem (Apter, J, Löwe)

Let κ be an inaccessible Suslin cardinal. Then
 $(\kappa, \kappa^+, \kappa^{++}) \rightarrow (\kappa, \kappa^+, \kappa^{++})^{\kappa, \kappa, \kappa}$.

This extends a result of **Kechris** for the countable exponent case.

Theorem

For all regular κ with $\delta_{2n+1}^1 < \kappa < \delta_{2n+3}^1$ we have $\kappa \rightarrow (\kappa)^{\delta_{2n+1}^1}$ but
 $\kappa \not\rightarrow (\kappa)^{\delta_{2n+2}^1}$.

Corollary

$\aleph_{\omega \cdot 2 + 1}$, $\aleph_{\omega^\omega + 1}$ are regular cardinals without the weak partition property.

Application

In [Apter, J, Löwe] we used the first theorem to force over models of $ZF + AD$ to change the cofinalities of $\aleph_1, \aleph_2, \aleph_3$.

- ▶ There are 3 possibilities for \aleph_1 Namely, $\text{cof} = \omega, r$ (regular, non-measurable), or m (measurable).
- ▶ There are 4 possibilities for \aleph_2 , and 5 possibilities for \aleph_3 .
- ▶ 13 of the 60 total possibilities are “trivially inconsistent.” For example, \aleph_1 regular, $\text{cof}(\aleph_2) = \aleph_1$, and $\text{cof}(\aleph_3) = \aleph_2$.

Theorem (Apter, J, Löwe)

Assuming suitable large cardinals, all of the remaining 47 cases are consistent with ZF.

Types of Suslin Cardinals

By a **Lévy** pointclass we mean a pointclass Γ closed under \exists^{ω^ω} or \forall^{ω^ω} (or both),

The Wadge hierarchy of Lévy pointclasses falls into **projective hierarchies** of 4 types.

We specialize to the Suslin pointclasses. The limit Suslin cardinals κ correspond to the bases of projective hierarchies.

Note that if κ is a limit Suslin cardinal then $\Lambda = S(< \kappa)$ is a selfdual pointclass closed under quantifiers.

Also, $o(\Lambda) = \delta(\Lambda) = \kappa$.

If $\text{cof}(\kappa) > \omega$, there is a nonselfdual pointclass Γ_S (Steel pointclass) of Wadge degree κ closed under \forall^{ω^ω} , \wedge , with $\text{sep}(\check{\Gamma}_S)$. Also, $\text{scale}(\Gamma_S)$.

Fact (Steel)

If κ is regular, then Γ_S is closed under \vee, \wedge .

When κ is regular, we have boundedness of Δ_S sets with respect to Γ_S -norms on Γ_S -complete sets.

- ▶ **Type I** $\text{cof}(\kappa) = \omega$. Let $\Sigma_0^\kappa = \bigcup_\omega (S(< \kappa))$. Then $\text{scale}(\Sigma_0^\kappa)$, $\text{scale}(\Pi_1^\kappa)$, $\text{scale}(\Sigma_2^\kappa), \dots$. The Suslin cardinals are $\kappa = \lambda_1^\kappa$, $\kappa^+ = \delta_1^\kappa, \lambda_3^\kappa, \delta_3^\kappa, \dots$
- ▶ **Type II** $\text{cof}(\kappa) > \omega$, Γ_S not closed under \vee .
- ▶ **Type III** $\text{cof}(\kappa) > \omega$, Γ_S closed under \vee but not $\exists^{\omega^\omega} \kappa$ necessarily regular).
- ▶ **Type IV** $\text{cof}(\kappa) > \omega$, Γ closed under $\exists^{\omega^\omega}, \forall^{\omega^\omega}$.

For κ an inaccessible Suslin cardinal we are in Type III or Type IV.

The Type I hierarchies will play an important role in the proof.


We fix the inaccessible Suslin cardinal κ . Let $C \subseteq \kappa$ be the c.u.b. set of limit Suslin cardinals. Let $C_\omega \subseteq C$ be the points of cofinality ω .

κ has the strong partition property by Kechris-Kleinberg-Moschovakis-Woodin.

Let μ be the ω -cofinal, normal measure on κ .

For $\alpha \in C_\omega$, let μ_α be the ω -cofinal, normal measure on α^+ . We have $j_{\mu_\alpha}(\alpha^+) = \alpha^{++}$.

We have the following picture.



$$\Sigma_0^\alpha = \bigcup_{\lambda < \alpha} S(\lambda)$$

$$S(\alpha) = \Sigma_1^\alpha, \quad S(\alpha^+) = \Sigma_2^\alpha$$

$$\text{scale}(\Sigma_0^\alpha,) \quad \text{scale}(\Pi_1^\alpha)$$

Plan of the proof

1. We show $[\alpha \mapsto \alpha^+]_\mu = \kappa^+$.
2. We show $\delta \doteq [\alpha \mapsto \alpha^{++}]_\mu \leq \kappa^{++}$.
3. We show for all $\theta < \omega_1$ that $(\kappa, \kappa^+, \delta) \rightarrow (\kappa, \kappa^+, \delta)^\theta$.
It follows that δ is regular, so $\delta = \kappa^{++}$.
4. Finally, we show $(\kappa, \kappa^+, \delta) \rightarrow (\kappa, \kappa^+, \delta)^\kappa$

The Trees T^+ and T^{++}

We define two trees T^+ and T^{++} on $\omega \times \kappa$.

Lemma

For any $f: \kappa \rightarrow \kappa$ with $f(\alpha) < \alpha^+$ there is a $x \in \omega^\omega$ with T_x^+ wellfounded and such that $\forall_\mu^ \alpha \ f(\alpha) < |T_x^+ \upharpoonright \alpha|$.*

Lemma

For any $f: \kappa \rightarrow \kappa$ with $f(\alpha) < \alpha^{++}$ there is a $x \in \omega^\omega$ with T_x^{++} wellfounded such that $\forall_\mu^ \alpha \ (f(\alpha) < [\beta \mapsto |T_x^{++} \upharpoonright \beta|]_{\mu_\alpha})$.*

The first lemma shows $[\alpha \mapsto \alpha^+]_\mu \leq \kappa^+$, and it follows easily that $[\alpha \mapsto \alpha^+]_\mu = \kappa^+$.

The second lemma shows that $\delta \doteq [\alpha \mapsto \alpha^{++}]_{\mu_\square} \leq \kappa^{++}$.

Construction of T^+

Fix a Γ -complete set P and a Γ -scale $\{\varphi_n\}_{n \in \omega}$ on P . we use $\varphi = \varphi_0$ to code ordinals $< \kappa$.

Say α is **strongly reliable** if for all $\beta < \alpha$:

$$\sup\{\varphi_n(x) : x \in P \wedge \varphi_0(x) \leq \beta\} < \alpha$$

The set of strongly reliable ordinals is c.u.b. in κ . Assume $C \subseteq$ strongly reliables.

Let

$$(x, y) \in R \leftrightarrow (x, y \in P \wedge \varphi(x) < \varphi(y)).$$

and

$$(x, y) \in R^\alpha \leftrightarrow (x, y \in P \wedge \varphi(x) < \varphi(y) < \alpha).$$

$R^\alpha \in \Sigma_0^\alpha - \bigcup_{\lambda < \alpha} \mathcal{S}(\lambda)$, and we uniformly have a Σ_0^α scale on R^α (essentially by restricting the scale $\vec{\varphi}$ to ordinals below α).

Starting from this, we uniformly get Σ_1^α universal sets B^α and Π_1^α sets Q^α and Π_1^α scales $\vec{\psi}^\alpha$ on Q^α .

Definition

Let $W = \{x : \forall n (x)_n \in P\}$. $x \in W$ will code the ordinal $|x| = \sup_n \varphi_0((x)_n)$.

The scale on P easily gives a scale on W . Let T_W be the corresponding tree.

We first construct a tree U on $\omega \times \omega \times \kappa$ with the following properties:

1. If $x \in W$ and $|x| = \alpha \in \mathcal{C}$, then $U_{x,y}$ is wellfounded iff the Σ_1^α relation coded by y is wellfounded.
2. For x, y as above, $|U_{x,y} \upharpoonright \alpha| \geq |B_y^\alpha|$, the Σ_1^α relation coded by y .

Key Point: For x, y as above, the entire tree $U_{x,y}$ is wellfounded (not just $U_{x,y} \upharpoonright \alpha$).

idea: U is constructed as in the proof of the Kunen-Martin theorem, but we use the components of the real x to verify the appropriate reals are in $B_y^{|x|}$.

Suppose $f: \kappa \rightarrow \kappa$ and $f(\alpha) < \alpha^+$ for $\alpha \in C$.

Consider the game G_f :

- I r
- II x, y

II wins the run iff

$$(r \in W) \rightarrow (x \in W \wedge B_y^{|x|} \text{ is wellfounded} \wedge |B_y^{|x|}| > f(|x|).$$

A boundedness argument shows that II has a winning strategy.

This suggests the following definition of the tree T^+ on $(\omega)^2 \times \kappa \times (\omega)^2 \times \kappa$:

$(\sigma, r, \vec{\alpha}, x, y, \vec{\beta}) \in [T^+]$ iff:

1. $(r, \vec{\alpha}) \in [T_W]$.
2. $\sigma(r) = (x, y)$
3. $(x, y, \vec{\beta}) \in [U]$.

Then T_σ^+ is wellfounded and $|T_\sigma^+ \upharpoonright \alpha| > f(\alpha)$ for μ almost all α .

Construction of T^{++}

We first construct a tree V on $(\omega)^2 \times \kappa$ with the following properties:

1. $(x, y) \in [V]$ iff $x \in W$ and for all n , $(y)_n$ codes a $\Sigma_1^{|x|}$ wellfounded relation $B_{(y)_n}^{|x|}$.
2. If $x \in W$, $|x| \in \mathcal{C}$ then there is a c.u.b. $D \subseteq \alpha^+$ such that if $\gamma \in D$, $y \in \omega^\omega$ and for all n $(y)_n$ codes a $\Sigma_1^{|x|}$ wellfounded relation of rank $< \gamma$, then $V_{x,y} \upharpoonright \gamma$ is illfounded.

main point: We can translate the $\Pi_1^{|x|}$ statement asserting the wellfoundedness of the $B_{(y)_n}^{|x|}$ into Π_1^β statements for any $\beta \geq |x|$ (use the $(x)_i$ as in the definition of U).

Suppose $x \in W$, $|x| = \alpha \in C$, and $g: \alpha^+ \rightarrow \alpha^+$. Play the game G_g :

I z
II w

II wins the run iff:

$(\forall n B_{(z)_n}^\alpha \text{ is wellfounded}) \rightarrow (B_w^\alpha \text{ is wellfounded} \wedge |B_w^\alpha| > g(\sup_n |B_{(y)_n}^\alpha|))$

By boundedness, II has a winning strategy τ for any G_g .

Suppose now $f: \kappa \rightarrow \kappa$ with $f(\alpha) < \alpha^{++}$.

Play the game G_f :

I r
II x, τ

r, x will be in W and τ will be strategy for a game G_g where $[g]_{\mu_\alpha} > f(\alpha)$, where $\alpha = |x|$.

More precisely, II wins the run iff:

$$\begin{aligned} r \in W &\rightarrow (x \in W \wedge |x| = \alpha \geq |y|) \\ &\wedge \forall z [\forall n B_{(z)_n}^\alpha \text{ is wellfounded} \rightarrow \\ &\quad B_{\tau(z)}^\alpha \text{ is wellfounded} \wedge |B_{\tau(z)}^\alpha| \geq g(\sup_n |B_{(z)_n}^\alpha|)] \end{aligned}$$

for some $g: \alpha^+ \rightarrow \alpha^+$ with $[g]_{\mu_\alpha} \geq f(\alpha)$.

II has a winning strategy σ for any f , and this suggests the definition of T^{++} :

$(\sigma, r, \vec{\alpha}, x, \tau, y, z, \vec{\beta}, \vec{\gamma}) \in T^{++}$ iff:

1. $(r, \vec{\alpha}) \in [T_W]$.
2. $\sigma(r) = (x, \tau)$.
3. $(x, y, \vec{\beta}) \in [V]$
4. $\tau(y) = z$.
5. $(x, a, \vec{\gamma}) \in [U]$

The properties of U and V show that T^{++} has the desired property.

The countable exponent θ case.

Fix a bijection $\pi: \omega \cdot \theta \rightarrow \omega$.

We code cofinally in κ^+ , κ^{++} many ordinals using sections of our trees: T_x^+ , T_x^{++} .

Suppose \mathcal{P} is a partition of the block functions from $3 \times \theta$ to $(\kappa, \kappa^+, \kappa^{++})$.

Consider the game $G_{\mathcal{P}}$:

- I x, y, z
- II x', y', z'

(1) If there is an $j < \omega \cdot \theta$ such that $(x)_{\pi(j)} \notin P_0$ or $(x')_{\pi(j)} \notin P_0$, then player I wins iff for the least such j , $(x)_{\pi(j)} \in P_0$.

(2) Suppose next that there is an $\alpha < \kappa$ such that one of the following holds.

(a) There is a $j < \omega \cdot \theta$ such that either $T_{(y)_{\pi(j)}}^+ \upharpoonright \alpha$ or $T_{(y')_{\pi(j)}}^+ \upharpoonright \alpha$ is illfounded.

(b) There is a $\beta < \alpha^+$ and a $j < \omega \cdot \theta$ such that either $T_{(z)_{\pi(j)}}^{++} \upharpoonright \beta$ or $T_{(z')_{\pi(j)}}^{++} \upharpoonright \beta$ is illfounded.

Let $\alpha < \kappa$ be least such that (a) or (b) above holds. If (a) holds, let j be least such that (a) holds for α and this j . In this case, Player I wins provided $T_{(y)_{\pi(j)}}^+$ is wellfounded. If (a) does not hold at α , but (b) does, let (β, j) be lexicographically least such that (b) holds. Player I wins in this case provided $T_{(z)_{\pi(j)}}^{++} \upharpoonright \beta$ is wellfounded.

Assume II has a winning strategy τ .

We define c.u.b. sets $C_0 \subseteq \kappa$, $C_1 \subseteq \kappa^+$, and $C_2 \subseteq \kappa^{++}$.

For example, to define C_2 we define for $\alpha \in C$, $\beta, \gamma < \alpha^+$ and $j < \omega \cdot \theta$:

$$A_{\alpha, \beta, \gamma, j} = \{(x, y, z) ; \forall j ((x)_{\pi(j)} \in P_0 \wedge \varphi_0((x)_{\pi(j)}) < \alpha) \\
 \wedge \forall \alpha' < \alpha \forall \beta < (\alpha')^+ \forall j (T_{(y)_{\pi(j)}}^+ \upharpoonright \alpha \text{ and } T_{(z)_{\pi(j)}}^{++} \upharpoonright \beta \text{ are wellfoun} \\
 \wedge \forall j |T_{(y)_{\pi(j)}}^+ \upharpoonright \alpha| < \beta \wedge \forall (\beta', j') \leq_{\text{lex}} (\beta, j) (|T_{(z)_{\pi(j)}}^{++} \upharpoonright \beta| \leq \eta)\}.$$

We have: $A_{\alpha,\beta,\gamma,j} \in \Delta_1^\alpha$.

Since τ is winning for Player II, for each $(x, y, z) \in A_{\alpha,\beta,\eta,j}$, if $\tau(x, y, x) = (x', y', z')$ then $\forall (\beta', j') \leq_{\text{lex}} (\beta, j) T_{(z')\pi(j')}^{++} \upharpoonright \beta$ is wellfounded.

By boundedness,

$$\rho_2(\alpha, \beta, \eta, j) := \sup\{|T_{(z')\pi(j')}^{++} \upharpoonright \beta|; (x', y', z') \in \tau[A_{\alpha,\beta,\eta,j}] \wedge j' \leq j\} < \alpha^+.$$

Let $C_2^\alpha \subseteq \alpha^+$ be c.u.b. closed under ρ_2 . The C_2^α lift to $C_2 \subseteq \kappa^{++}$.

Exponent κ

We use generic codes (**Kechris-Woodin**) and the uniform coding lemma.

Let $U(Q, z, x, y)$ be universal for the syntactic class $\Sigma_1(Q)$, where Q is a binary predicate symbol.

Can take

$$U(Q, z, x, y) \leftrightarrow \exists w (S(z, \langle x, y, w \rangle) \wedge \forall n W(\langle (w)_n \rangle_0, \langle (w)_n \rangle_1)).$$

where S is universal Σ_1^1 .

Let R'_α code $\{(x, y) : \varphi(x) < \varphi(y) \leq \alpha\}$ and \leq version.

We write $U_z(R'_\alpha)(x, y)$ for $U(R'_\alpha, z, x, y)$.

Uniform Coding Lemma says that if $A \subseteq \omega^\omega \times \omega^\omega$ with $\text{dom}(A) = P$, then there is a $z \in \omega^\omega$ such that for all $\alpha < \kappa$, $U_z(R'_\alpha)$ is a choice subrelation for $A \upharpoonright P_{\leq \alpha}$.

Let $\overline{U}_z(R'_\alpha)$ be a uniformization of $U_z(R'_\alpha)$ (using scale $\{\varphi_n\}$).

For $\alpha < \kappa$, let α' be the next reliable (w.r.t. $\{\varphi_n\}$).

Let $G: \kappa^\omega \rightarrow \omega^\omega$ be a **generic coding function**. So for any $s \in \kappa^\omega$, $x = G(\alpha \hat{\ } s) \in P$, $\varphi(x) \leq \alpha$, and if s enumerates an honest set then $|x| = \alpha$.

Given reals x, y, z we say:

1. x **codes a function** at $\alpha < \kappa$ if $|a| = \alpha$ implies $\exists b \in P \overline{U}_x(R'_\alpha)(a, b)$. Also, if $|a| = |a'|$ then $|b| = |b'|$.
2. y is **good** at $\delta < \alpha \in C_\omega$ if $\forall^* a \in P_\delta \exists b \overline{U}_y(R'_\delta)(a, b)$ and $T_b^+ \upharpoonright \alpha$ is wellfounded (good at α if good at all $\delta < \alpha$).
3. z is **good** at $\delta < \alpha \in C_\omega, \beta < \alpha^+$ if $\forall^* a \in P_\delta \exists b \overline{U}_z(R'_\delta)(a, b)$ and $T_z^{++} \upharpoonright \beta$ is wellfounded (good at α if good at all $\delta < \alpha, \beta < \alpha^+$).

$f_x(\alpha) =$ the unique value of $|b|$.

$g_y(\delta, \alpha) =$ least $\gamma < \alpha^+$ such that $\forall^* a \in P_\delta, |T_b^+ \upharpoonright \alpha| \leq \gamma$.

$h_z(\delta, \alpha, \beta) =$ least $\gamma < \alpha^+$ such that $\forall^* a \in P_\delta, |T_b^{++} \upharpoonright \beta| \leq \gamma$.

I plays x, y, z , II plays x', y', z' .

Let α be least such that one of the following holds.

1. For some $\delta < \alpha$, y or y' is not good at (δ, α) .
2. For some $\delta < \alpha, \beta < \alpha^+$, z or z' is not good at (δ, α, β) .
3. x or x' does not code a function at α .

If (1) holds, then I wins iff for the least such δ , y is good at (δ, α) .
Suppose (1) does not hold, but (2) holds. Then I wins iff for the lexicographically least pair (β, δ) we have z is good at δ, α, β . If (1) and (2) don't hold, but (3) holds, then I wins iff x codes an ordinal at α .

Otherwise, we have functions $f_x, f_{x'}, g_y, g_{y'}, h_z, h_{z'}$.

- ▶ $f_x, f_{x'}$ together produce $F: \kappa \rightarrow \kappa$.
- ▶ $g_y, g_{y'}$ together produce $G: \kappa \rightarrow \kappa^+$.
- ▶ $h_z, h_{z'}$ together produce $H: \kappa \rightarrow \kappa^{++}$.

II wins the run in this case iff $\mathcal{P}(F, G, H) = 1$.

From a winning strategy τ for II, say, we define the homogeneous sets $C_0 \subseteq \kappa$, $C_1 \subseteq \kappa^+$ and $C_2 \subseteq \kappa^{++}$.